

UNCLASSIFIED

AD NUMBER

AD861786

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors;
Administrative/Operational Use; 31 DEC 1986.
Other requests shall be referred to Air Force Technical Applications Center, Washington, DC.

AUTHORITY

AFTAC USAF ltr 25 Jan 1972

THIS PAGE IS UNCLASSIFIED

AD 861786

RAYLEIGH WAVE ANALYSIS OF ATMOSPHERIC EXPLOSIONS

31 December 1968

Prepared For


AIR FORCE TECHNICAL APPLICATIONS CENTER
Washington, D. C.

By

L. S. Turnbull
D. B. Rabenstine
TELEDYNE, INC.

Under

Project VELA UNIFORM



Sponsored By

ADVANCED RESEARCH PROJECTS AGENCY
Nuclear Test Detection Office
ARPA Order No. 624

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Chief, AFTAC.

attn: VSC, Wash D.C. 20333

BLANK PAGE

RAYLEIGH WAVE ANALYSIS OF ATMOSPHERIC EXPLOSIONS

SEISMIC DATA LABORATORY REPORT NO. 228

AFTAC Project No.:	VELA T/6702
Project Title:	Seismic Data Laboratory
ARPA Order No.:	624
ARPA Program Code No.:	8F10
Name of Contractor:	TELEDYNE INDUSTRIES, INC.
Contract No.:	F 33657-68-C-0945
Date of Contract:	2 March 1968
Amount of Contract:	\$ 1,251,000
Contract Expiration Date:	1 March 1969
Project Manager:	Royal A. Hartenberger (703) 836-7647

P. O. Box 334, Alexandria, Virginia

AVAILABILITY

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Chief, AFTAC.

This research was supported by the Advanced Research Projects Agency, Nuclear Test Detection Office, under Project VELA-UNIFORM and accomplished under technical direction of the Air Force Technical Applications Center under Contract F 33657-68-C-0945.

Neither the Advanced Research Projects Agency nor the Air Force Technical Applications Center will be responsible for information contained herein which may have been supplied by other organizations or contractors, and this document is subject to later revision as may be necessary.

ABSTRACT

During the months of July through September, 1968, a series of atmospheric nuclear explosions was detonated by France in the South Pacific. The events analyzed are those of July 7, July 15, August 3, August 24, and September 3, 1968. The analysis includes matched filtering for relative amplitude (\hat{a}) measurements, body and surface wave magnitude comparisons phased summing, and ARZ-ERZ measurements. The events are discussed individually, and a summary of pertinent data is presented in several figures and tables.

TABLE OF CONTENTS

	Page No.
ABSTRACT	
INTRODUCTION	1
Data Analysis	1
A. July 7	3
B. July 15	3
C. August 3	4
D. August 24	5
E. September 8	5
CONCLUSIONS	7
REFERENCES	8

LIST OF FIGURES

Figure Title	Figure No.
Event of 7 July 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam formed on 3.75 km/sec. Visible event is the Philippine event described in the text	1
Event of 7 July 1968. LASA & LRSM LPZ Rayleigh waves, matched filtered with signals from the event of 24 August 1968. Interfering event is the Philippine event described in the text.	2
Event of 15 July 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.	3
Event of 15 July 1968. LASA LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.	4
Event of 15 July 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.	5
Event of 3 August 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.	6
Event of 3 August 1968. LASA LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.	7
Event of 3 August 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.	8
Event of 24 August 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.	9
Event of 8 September 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.	10

LIST OF FIGURES (Cont'd.)

Figure Title	Figure No.
Event of 8 September 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.	10
Event of 8 September 1968. LASA LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.	11
Event of 8 September 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.	12
M_s vs m_b comparison of various seismic signals.	13
ARZ vs m_b comparison of various seismic signals.	14
ERZ vs m_b comparison of various seismic signals	15

TABLE

Table Title	Table No.
Summary of \hat{a} Values	I

INTRODUCTION

During the months of July through September, 1968 a series of atmospheric nuclear explosions was detonated by France in the South Pacific. The events from that series analyzed in the report are those of July 7, July 15, August 3, August 24, and September 8, 1968. The analysis includes matched filtering (Alexander and Rabenstine, 1967a, 1967b) for relative amplitude ~~(\hat{a})~~ measurements, body and surface wave magnitude comparisons (Lambert and Turnbull, 1968), phased summing, and ARZ-ERZ measurements (Turnbull and Lambert, 1968). The events will be discussed individually in the following sections, with a summary of pertinent data presented in several figures and tables. () <

Data Analysis

The five events of interest occurred essentially at the same coordinates. The data we wished to analyze were those recorded at LASA, TFO, UBO, and five LRSM stations (LCNM, MNNV, PGBC, RKON, and SV3QB). The data from TFO and UBO proved unusable because neither LP array was yet operational.

The LASA data were recorded digitally at 5 samples per second and decimated to 1 sample per second for analysis. The LRSM data were recorded on analog tape, digitized at the SDL at 5 samples per second, and then decimated to 1 sample per second for analysis.

The LASA data were demagnified using results of .04 Hz sine wave electro-magnetic calibrations recorded in March 1967.

Obviously these calibrations are not accurate for current data, but they are the most recent that we have available. This, however, should not affect the relative amplitude (\hat{a}) values. The LRSM data were demagnified by comparison with the film records and indicated magnifications.

Only the sensors in the A,D,E, and F subarrays at LASA were used, in order to assure incoherent noise (Alexander and Rabenstine, 1967b).

The primary technique of our analysis was the matched filter. A search for events which originated in the same source region with sufficient magnitude to generate a good dispersed wavetrain was undertaken using the library program DEPTHMAG. The source region was shown to be extremely aseismic, with only one event meeting the needed specifications (i.e., within 10°). An earthquake occurred at 18.4S, 132.9W of magnitude 5.5 on March 6, 1965. LASA was not in operation at that time. However, an ideal matched filter was furnished by one of the explosions itself; the event of August 24. The body wave magnitude was 4.8 and the surface wave magnitude was found to be 4.7. A dispersed wavetrain with good S/N was readily visible on the records at LASA. Thus, the \hat{a} values measured are amplitudes of the various events relative to the event of August 24.

The bandpass filter which was used on the raw data either before phased summing or when matched filtering was simply a shaping applied to the Fourier transform. The transform was multiplied by a linear taper which went from 1.0 to 0.0 as the frequency went from 0.02 to 0.0167 Hz and 0.1 to 0.125 Hz. Transform values outside the frequency range of 0.0167 to 0.125 Hz were set to zero.

A. July 7

Long period analysis of this event by all techniques proved quite futile. Approximately one-half hour before detonation, an earthquake of magnitude 5.1* (m_b) occurred in Mindanao, Phillipines (9.6N, 126.5E). The paths were such that the wavetrains of the earthquake and the explosion overlapped at LASA and most LRSM stations. Results of the phased sum are shown in Figure 1 and of the matched filter and phased sum in Figure 2.

B. July 15

This event was determined to have a body wave magnitude (m_b) of 4.3* and a surface wave magnitude (M_s) of 4.1. The surface wave magnitude is considerably larger than that for an underground nuclear explosion of comparable body wave magnitude (approximately two orders of magnitude), an indication that an atmospheric nuclear explosion generates a great deal more long period energy. In a plot of M_s vs. m_b (Figure 13), the events of this series are shown along with earthquakes and underground nuclear explosions. For the same order body wave magnitude, we find that atmospheric explosions generate even more long period energy than earthquakes. The energy difference is further displayed in plots of ARZ vs. m_b (Figure 14) and ERZ vs. m_b (Figure 15).

The signal as recorded at LASA was readily visible (Figure 3). The records of eleven long-period instruments

* All body wave magnitudes are from the LASA bulletin; all surface wave magnitudes are computed in the ARZ-ERZ program (Turnbull and Lambert, 1968).

were band-pass filtered from 10 to 50 seconds (period) and phase summed using a velocity of 3.75 km/sec. From an average S/N ratio of 4.0 for each channel, the sum channel S/N was 9.5.

The LASA matched filter output was quite impressive, as could be expected from the visible signal (Figure 4). The S/N ratio of the matched filter output had an average of 11.2 (\hat{a} avg. = 0.248), and the S/N ratio of the sum of the matched filter outputs from eleven channels was 34.7 (\hat{a} = 0.238). The summed matched filter output was quite symmetric and narrow indicating high correlation between the filter and event. The standard deviation of the \hat{a} values was 14% of the mean value.

When the five LRSM signals were matched filtered and beam formed with the LASA matched filter outputs on 4 km/sec, a further S/N enhancement to 47.1 was achieved (Figure 5), with an \hat{a} of 0.209 on the sum. The mean of all \hat{a} values was 0.234 with a standard deviation of 16%.

C. August 3

This event was determined to have a body wave magnitude (m_b) of 4.1 and a surface wave magnitude (M_s) of 4.3. As in the July 15 event, a great deal more long period energy than in an underground explosion of the same m_b is generated. However, the signal as recorded at LASA was not as discernible as the July 15 event (Figure 6), the average S/N being 3.1 and the S/N of the phase summed trace of eleven channels only 3.8.

This reason for the very small indicated increase in S/N is probably that the signal levels measured on the individual traces were too high because of noise being mistakenly measured as signal.

The LASA matched filter output, though, showed a greater improvement upon summation (Figure 7). While the average S/N for eleven matched filtered channels was still only 3.1 (\hat{a} avg. = 0.0695), the sum S/N was 8.3 (\hat{a} = 0.06). The standard deviation of \hat{a} values was 32%, again due to the relatively low S/N ratios.

Once again beamforming the 5 LRSM matched filtered signals with the LASA matched filtered signals on a velocity of 4 km/sec increased the S/N to 9.56 (Figure 8) with a sum trace \hat{a} of 0.057. The mean of all \hat{a} values was 0.0698 with a standard deviation of 24%.

D. August 24

This event had a body wave magnitude (m_b) of 4.8 and a surface wave magnitude (M_s) of 4.75. The Rayleigh-wave train was very well defined on the LASA records, hence its use as the matched filter for the other events (Figure 9). The S/N for an average band pass filtered channel was 17.9, while the S/N of the phased sum of thirteen channels is 51.7.

E. September 8

This event was the second largest of the series and had a body wave magnitude of 4.6. The surface wave magnitude was measured to be 4.6, and the signal at LASA was well recorded. The previous conclusions on surface wave energy from atmospheric explosions are again verified.

The beam forming of the band pass filtered surface waves at LASA worked quite well giving an improvement in S/N from a mean of 6.3 on the individual traces to 14.1 on the sum trace (Figure 10.).

The matched filter also worked well, as expected with a large signal. The LASA matched filtered traces achieved a mean S/N of 13.51 (\hat{a} avg. = 0.541) and summed to give a S/N of 37.77 (\hat{a} = 0.508) (Figure 11). The LRSM matched filtered data, which had mean S/N of 11.35 (\hat{a} avg. = 0.449), combined with the LASA data to give a sum S/N of 42.33 (\hat{a} = 0.488) (Figure 12). The standard deviation of all the \hat{a} values was 22% of the mean.

CONCLUSIONS

1. In comparing atmospheric explosions with underground explosions, we find the former generate more long period energy than the latter, with the surface wave magnitudes generally two orders of magnitude greater.

2. The "best" matched filter proved to be a large atmospheric shot from the same region.

3. The LRSM matched filtered data was beam formed with the LASA matched filtered data to give S/N values very near that which is predicted for coherent signals in incoherent noise. The exception was a case where the actual values of S/N on the individual LASA channels were too low to measure accurately.

4. Reasonably consistent values of \hat{a} were obtained across the network of stations used.

REFERENCES

- Alexander, S.S., and D.B. Rabenstine, 1967a, Detection of surface waves from small events at teleseismic distances, SDL Report No. 175.
- Alexander, S.S., and D.B. Rabenstine, 1967b, Rayleigh wave signal to noise enhancement for a small teleseism using LASA, LRSM, and observatory stations, SDL Report No. 194.
- Lambert, D.G. and L.S. Turnbull, 1968, Personal Communication.
- Turnbull, L.S. and D.G. Lambert, 1968, Rayleigh wave discrimination techniques between underground explosions and earthquakes, SDL Report No. 211.

Table I Summary of \hat{a} Values

	15 JULY			3 AUGUST			8 SEPTEMBER		
<u>LASA</u>	\hat{a}	(S/N)	n	\hat{a}	(S/N)	n	\hat{a}	(S/N)	n
MEAN	<u>.2475</u>	(11.20)**	11	<u>.0695</u>	(3.13)**	11	<u>.5407</u>	(13.51)**	12
MAX	.3021	(7.63)*		.1007	(4.99)*		.8482	(14.55)*	
MIN	.1758	(12.56)*		.0361	(1.47)*		.3191	(16.31)*	
SQ. MEAN	.061256			.0048289			.292356		
MEAN SQ.	.062469			.0051970			.307805		
VARIANCE	.001213			.0003681			.015449		
STD. DEV	<u>.0348</u>			<u>.0192</u>			<u>.1243</u>		
SUM VAL	<u>.2379</u>	(34.68)*		<u>.0605</u>	(8.33)*		<u>.5075</u>	(37.77)*	
<u>LRSM</u>									
MEAN	<u>.2041</u>	(15.76)**	5	<u>.0706</u>	(2.53)**	5	<u>.4485</u>	(11.35)**	5
MAX	.2255	(16.44)*		.0820	(2.73)		.4862	(17.46)*	
MIN	.1467	(13.95)*		.0566	(1.86)*		.3779	(8.25)*	
SQ. MEAN	.041657			.0049855			.201152		
MEAN SQ.	.042536			.0050672			.202810		
VARIANCE	.000873			.0000817			.001658		
STD. DEV	<u>.0296</u>			<u>.0090</u>			<u>.0407</u>		
SUM VAL	<u>.1620</u>	(34.30)*		<u>.0480</u>	(4.06)*		.4355	(29.66)*	
<u>ALL</u>									
MEAN	.2340	(12.63)**	16	.0698	(2.94)**	16	.5136	(12.88)**	17
MAX	.3021	(7.63)*		.1007	(4.99)*		.8482	(14.55)*	
MIN	.1467	(13.95)*		.0361	(1.47)*		.3191	(16.31)*	
SQ. MEAN	.054756			.0048776			.263785		
MEAN SQ.	.056240			.0051564			.276924		
VARIANCE	.001484			.0002788			.013139		
STD. DEV	<u>.0385</u>			<u>.0167</u>			<u>.1146</u>		
SUM VAL	<u>.2085</u>	(47.08)*		<u>.0570</u>	(3.56)*		<u>.4879</u>	(42.33)*	

* S/N of trace with with indicated \hat{a}
 ** Mean S/N of traces in set
 n = number of traces in set




Figure 1. Event of 7 July 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam formed on 3.75 km/sec. Visible event is the Philippine event described in the text.




Figure 2. Event of 7 July 1968. LASA & LRSM LPZ Rayleigh waves, matched filtered with signals from the event of 24 August 1968. Interfering event is the Philippine event described in the text.




Figure 3. Event of 15 July 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.




Figure 4. Event of 15 July 1968. LASA LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.




Figure 5. Event of 15 July 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.




Figure 6. Event of 3 August 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.




Figure 7. Event of 3 August 1968. LASA LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.




Figure 8. Event of 3 August 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.




Figure 9. Event of 24 August 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.




Figure 10. Event of 8 September 1968. LASA LPZ Rayleigh waves, band pass filtered (10-50 sec. period) and beam-formed on 3.75 km/sec.




Figure 11. Event of 8 September 1968. LASA LPZ Rayleigh waves matched filtered with signals from event of 24 August 1968. Sum of LASA matched filtered signals.




Figure 12. Event of 8 September 1968. LRSM LPZ Rayleigh waves, matched filtered with signals from event of 24 August 1968. Sum of LASA and LRSM matched filtered signals.




Figure 13. M_s vs m_b comparison of various seismic signals




Figure 14. ARZ vs m_b comparison of various seismic signals




Figure 15. ERZ vs m_b comparison of various seismic signals

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

TELEDYNE INDUSTRIES, INC.

ALEXANDRIA, VIRGINIA

2a. REPORT SECURITY CLASSIFICATION

Unclassified

2b. GROUP

3. REPORT TITLE

RAYLEIGH WAVE ANALYSIS OF ATMOSPHERIC EXPLOSIONS

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Scientific

5. AUTHOR(S) (Last name, first name, initial)

Turnbull, L.S.; Rabenstine, D.B.

6. REPORT DATE

31 December 1968

7a. TOTAL NO. OF PAGES

26

7b. NO. OF REFS

4

8a. CONTRACT OR GRANT NO.

F 33657-68-C-0945

9a. ORIGINATOR'S REPORT NUMBER(S)

228

8b. PROJECT NO.

VELA T/6702

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

ARPA Order No. 624

ARPA Program Code No. 8F10

10. AVAILABILITY/LIMITATION NOTICES

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Chief, AFTAC.

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

ADVANCED RESEARCH PROJECTS AGENCY
NUCLEAR TEST DETECTION OFFICE
WASHINGTON, D. C.

13. ABSTRACT

During the months of July through September, 1968, a series of atmospheric explosions was detonated in the South Pacific. The events analyzed are those of July 7, July 15, August 3, August 24, and September 3, 1968. The analysis includes matched filtering for relative amplitude (a) measurements, body and surface wave magnitude comparisons phased summing, and ARZ-ERZ measurements. The events are discussed individually, and a summary of pertinent data is presented in several figures and tables.

14. KEY WORDS

Matched filters
beamforming
atmospheric explosions

magnitude discriminant
LASA
LRSM

Unclassified

Security Classification